

# Quintox<sup>®</sup> RAT & MOUSE BAIT

## A Stop-Feed Rodenticide

QUINTOX Rat and Mouse Bait contains the active ingredient, Cholecalciferol, better known as Vitamin D<sub>3</sub>, which takes a completely different biological pathway to killing rodents, including anticoagulant-resistant rodents.

Unlike anticoagulants, once a rodent eats a lethal dose of QUINTOX, all feeding stops. QUINTOX acts faster than anticoagulants, causing death in 2 to 4 days.

Rodents can consume a lethal dose of QUINTOX in a single day's feeding or accumulate it in smaller, multiple feedings over a period of days. Just 1/10 oz. (2.8 grams) of QUINTOX can kill a mouse, while 1/4 oz. (7 grams) is a lethal dose for most rats. Bait shyness is not a problem with QUINTOX because toxic symptoms do not start until after a lethal dose is consumed.

QUINTOX may be used in and around homes, farms and commercial locations and is authorized by USDA for use in federally inspected meat and poultry plants.

**Packaging:** 5½ lb. bulk plastic pails – EPA REG. NO. 12455-39  
50 x 30 g. place pacs – EPA REG. NO. 3240-42-12455



## Specimen Label

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions.

**IMPORTANT:** Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused, spoiled, and unconsumed bait as specified on this label.

**USE RESTRICTIONS:** For control of Norway rats, roof rats and house mice in and around homes, industrial, commercial and agricultural buildings, and similar man-made structures. This product may also be used inside transport vehicles (ships, trains, aircraft) and in related port or terminal buildings.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats and mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

### APPLICATION DIRECTIONS

**RATS:** Apply 2 to 8 ounces of bait per placement. Space placements at intervals of 15 to 30 feet. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of rat activity cease.

**MICE:** Apply 1/4 to 1/2 oz. (1-2 level tablespoons) of bait at each placement location. Space placements at 8- to 12-foot intervals. Larger placements (up to 1 ounce) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of mouse activity cease.

**Quintox<sup>®</sup>**  
**RAT & MOUSE BAIT**

**KILLS RATS AND MICE**

ACTIVE INGREDIENT: Cholecalciferol ..... 0.075%  
INERT INGREDIENTS..... 99.925%  
TOTAL 100.000%


### CAUTION

**KEEP OUT OF REACH OF CHILDREN**

(See right side panel for additional precautionary statements.)

For 24-hour emergency medical advice, call 1-877-854-2494.

Manufactured by:

 **Bell Laboratories, Inc.**  
Madison, WI 53704 U.S.A.

EPA EST. NO. 12455-WI-1 EPA REG. NO. 12455-39

**DIRECTIONS FOR USE** (Continued from other panel)

**APPLICATION DIRECTIONS** (Continued from other panel):  
**RATS AND MICE:** Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and left-over bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**

Keep away from humans, domestic animals, and pets.

### STATEMENT OF PRACTICAL TREATMENT

If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Avoid use of all oils.

**NOTICE TO PHYSICIAN:** If serum calcium levels are elevated, treatment with Calcitonin is effective in reducing calcium to normal levels. Continue monitoring serum calcium and treat as necessary for hypercalcemia (Reference, AMA Drug Evaluations, Third Edition (1977), Chapter 16, pp. 248-251).

### ENVIRONMENTAL HAZARDS

Keep out of lakes, streams or ponds.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store only in original container in a dry area inaccessible to children and pets.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty pail. Then dispose of empty pail in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**WARRANTY:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.