

# GENERATION SPECIMEN LABEL

## PRECAUTIONARY STATEMENTS

### HAZARD TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Keep away from children, domestic animals and pets. May be harmful or fatal if swallowed or absorbed through the skin because this material may reduce the clotting ability of blood and cause bleeding.

Do not get in eyes, on skin or on clothing. Wash arms, hands and face with soap and water after applying and before eating and smoking.

**IF SWALLOWED CALL A PHYSICIAN AT ONCE.**

**NOTE TO PHYSICIAN OR VETERINARIAN:** This product may reduce the clotting ability of the blood and cause hemorrhaging. If poisoning occurs, intramuscular and oral administration of Vitamin K<sub>1</sub> is antidotal as indicated in bishydroxycoumarin overdoses. Repeated doses may need to be given for up to two weeks (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

## STORAGE AND DISPOSAL

**DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL. STORE IN ORIGINAL CONTAINER IN LOCKED STORAGE AREA**

**STORAGE:** Store in original container in a cool, dry place inaccessible to children and pets.

#### PESTICIDE DISPOSAL:

Wastes resulting from the use of this product should be securely wrapped in several layers of newspaper and then disposed of in a sanitary landfill or at an approved waste disposal facility.

#### CONTAINER DISPOSAL:

Do not reuse pail. Rinse thoroughly before discarding in trash.

**KEEP AWAY FROM FOODSTUFFS**

**WARRANTY:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Satisfaction Guaranteed or Your Money Back

EPA Reg. No. 7173-211

EPA Est. No.: 7173-WI-1



## READY-TO-USE

**KILLS NORWAY RATS, ROOF RATS, HOUSE MICE AND WARFARIN-RESISTANT HOUSE MICE**

**NORWAY RATS AND HOUSE MICE CAN CONSUME A LETHAL DOSE IN ONE FEEDING WITH FIRST DEAD RODENTS APPEARING FOUR OR FIVE DAYS AFTER TREATMENT BEGINS.**

#### Active Ingredient:

3-[3-(4'-bromo[1,1'-biphenyl]-4-yl)-1,2,3,4-tetrahydro-1-naphthylenyl]-4-hydroxy-2H-1-benzothiopyran-2-one*	0.0025%
Inert Ingredients	99.9975%
Total	100.0000%

\*Difethialone - Patented in the U.S.A.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

MAY BE HARMFUL OR FATAL IF SWALLOWED.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

MANUFACTURED BY:

**LIPHA<sup>®</sup>TECH**

LiphaTech Inc.  
3101 West Custer Ave.  
Milwaukee, WI 53209

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions.

**IMPORTANT:** Do not expose children, pets or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, unused, spoiled, and unconsumed bait as specified on this label.

**USE RESTRICTIONS:** For control of Norway rats, roof rats, and house mice.

**URBAN AREAS:** This product may be used in and around homes and industrial, commercial and public buildings. This product may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. All applications "around" buildings must be made in tamper-resistant bait stations placed along the walls or other external parts of buildings (e.g., doorways, ramps, loading docks) where rats or mice might seek to gain entrance.

**NON-URBAN AREAS:** This product may be used inside homes and agricultural buildings.

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Remove as much alternative food as possible.

## APPLICATION DIRECTIONS:

**RATS:** Place at least 3 to 16 packs per placement. Placements usually should be placed at intervals of 15 to 30 feet. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease.

**MICE:** Use one bait pack per placement. Two packs may be placed at points of very high mouse activity. Space placements at 8- to 12-foot intervals. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease.

**RATS AND MICE:** Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.